Roll	No.
Kon	140.

STATISTICS Time: 20 Minutes

# Intermediate Part-I, Class 11th (1stA 323- III)

PAPER: I Marks: 17

Note:

101-11-23

Code: 6185

You have four choices for each objective type question as A, B, C and D. The choice which you think is correct, fill that circle in front of that question number. Use marker or pen to fill the circles. Cutting or filling two more circles will result in zero mark in that question.

1-	1_	The standard	deviation	of hinomial	distribution	ic
1-	Ĭ -	THE Standard	deviation	OI UIIIUIIIIai	distribution	12

(A) np

- (B) npq
- (C)  $\sqrt{npq}$
- (D) nq

2- If 
$$\sum (X - \overline{X})^2 = 180$$
 and  $n = 9$ , then m<sub>2</sub> is

(A) 25

- (C) 20
- (D) 18

3- The index given by 
$$\frac{\sum p_n q_n}{\sum p_o q_n}$$
 is

- (A) Laspeyre's index
- (B) Paasche's index
- (C) Value index
- (D) Fisher index
- Let 'a' is a constant and 'X' is a random variable, then S.D.(aX) is 4-
  - (A)  $a^2$  S.D.(X)
- (B) S.D.(X)
- (C) aS.D.(X)
- (D) zero

- If  $B_2 = 3$ , then the distribution is called 5-
  - (A) mesokurtic
- (B) platykurtic
- (C) leptokurtic
- (D) ogive

- 6-Total angles of a pie chart are
  - (A)  $360^{\circ}$
- (B)  $180^{\circ}$
- (C) 190°
- (D) 90°

- The probability of an impossible event is equal to 7-
  - (A) zero

- (C) -1
- (D) 2

- The most frequent value in the data is 8-
  - (A) Mean
- (B) Median
- (C) Mode
- (D) G.M.

- Quantities which do not vary 9-
  - (A) variables
- (B) constants
- (C) statistics
- (D) all of these

- The Mean of Hypergeometric distribution is 10-

- (C)  $\frac{n}{N\kappa}$

- First moment about mean is always equal to 11-

(B) -1

- (C) 2
- (D) zero

- The number of parameters of binomial distribution, are 12-

(B) 3

- (C) 1
- (D) 4

- If two dice are rolled, the possible outcomes are 13-

(B) 12

- (C) 216
- (D) 36

- A single value which represents a distribution is called 14-
  - (A) S.D.
- (B) Variance
- (C) Average
- (D) C.V.

- Geometric mean of 2 and 8 is 15-

(B) 4

- (C) 5
- (D) 2

- In fixed base method the base period should be 16-

(A) far away

- (B) abnormal
- (C) normal
- (D) unreliable

- If  $f(x) = \frac{1}{10}$  and x=10, then E(X) is 17-
  - (A) zero
- (B) 1

- (C)  $\frac{1}{2}$
- (D) -1

**CATISTICS** 

Intermediate Part-I, Class 11th (1stA 323)

SUBJECTIVE (1

PAPER: I Marks: 68

Time: 2:40 Hours

Note: Section-I is compulsory. Attempt any Three (3) questions from Section-II.

SECTION-I

. Write short answers to any EIGHT (8) questions:

Cuj-11-23

 $(2 \times 8 = 16)$ 

- i- Define statistics.
- ii- Differentiate between variable and constant.
- iii- Write down any four qualities of a good average.
- iv- Define Geometric Mean.
- v- Given  $\sum (X-10) = 2.8$  and n = 5. Find Mean :  $\overline{X}$
- vi- Define weighted arithmetic mean.
- vii- Given L=60, h=10, f=20, n=80 and c=30. Find median.
- viii- Write down the empirical relationship between mean, median and mode.
- ix- Define Price Relative.
- x- Given  $P_{on}(Laspeyre's) = 120$ ,  $P_{on}(Paasche's) = 118$ . Find  $P_{on}(Fisher)$  price index number.
- xi- Given W=20, 25, 30, 40 and I=100, 105, 110, 120. Find consumer price index number.
- xii- What are the uses of index numbers?

#### 3. Write short answers to any EIGHT (8) questions:

 $(2 \times 8 = 16)$ 

- i- Define tabulation.
- ii- Define frequency distribution.
- iii- What is meant by absolute dispersion?
- iv- First, second and third quartiles of a distribution are 142, 153 and 167 respectively. Find coefficient of skewness.
- v- Write down two properties of variance.
- vi- What do you mean by kurtosis?
- vii- Given that n=8,  $\Sigma D = 10$ ,  $\Sigma D^2 = 524$ . Find variance, where D = X 15
- viii- Define mean deviation.
- ix- Define probability of an event.
- x- What are independent events?
- xi- State addition law of probability for mutually exclusive events.
- xii- State multiplication law of probability for two independent events.

### 4. Write short answers to any SIX (6) questions:

 $(2 \times 6 = 12)$ 

- i- What do you mean by probability density function?
- ii- Narrate two laws of expectation.
- iii- Given that E(x+4)=10 and  $E(x+4)^2=116$ . Find variance (x+4)
- iv- A continuous random variable X has probability density function

$$F(x) = c(4-x) \text{ for } 1 \le x \le 3$$

=0 elsewhere

Find the value of c.

- v- For a binomial distribution with n = 10 and  $p = \frac{1}{3}$ . Find P(X = 5)
- vi- If X is a hypergeometric random variable with N=8, n=6 and K=5. Find S.D.(X)
- vii- Describe hypergeometric experiment.
- viii- Write down the properties of a binomial experiment.
- ix- Describe hypergeometric probability distribution.

(Turn over)

0

## SECTION - II

Guj-11-23

### Note: Attempt any Three (3) questions.

5-	(a)	Find	the	G.M.	of the	following data
----	-----	------	-----	------	--------	----------------

Classes	10-19	20 – 29	30 – 39	40 – 49	50 - 59
f	5	25	40	20	10

(b) The reciprocals of 8 values of X are given below: 0.0500, 0.0454, 0.0400, 0.0333, 0.0285, 0.0232, 0.0213, 0.0200 Calculate A.M. and H.M.

6- (a) Find the coefficient of variation for the following data

Marks	1 – 3	3 – 5	5 – 7	7-9
f	10	15	20	25

(b) Find mean deviation from the following data

Group	2-4	4-6	6-8	8 – 10	10 – 12
f	3	45	6	4	3

7- (a) Compute Fisher Ideal index number using 2010 as base year, for the following data

	Pri	ces	Quar	tities
Commodities	2010	2015	2010	2015
A	10	12	120	100
В	8	10	150	130
C	12	13	80	70
D	15	20	60	50

- (b) A card is selected from a deck of playing cards. Find the probability that
  - i) The card is black
  - ii) The card is queen card
  - iii) The card is spade card
  - iv) The card is a face card

8- (a) A random variable 'X' has the following distribution

A faildoil variable X has the following distribution								
	X	0	1	2	3			
	P(X)	0.1	0.2	0.3	0.4			

Find (i) E(X) (ii) Var(X)

(b) Given the following probability distribution

)	Given the follow	ing probabilit	y distribution			
	Xi	0	1	2	3	4
	P(Xi)	1/126	20/126	60/126	40/126	5/126

Verify that E(2X+3)=2E(X)+3

- 9- (a) Out of 800 families with 5 children each, how many would you expect to have
  - i) At least 3 boys
  - ii) At most 1 boy
  - (b) Four balls are drawn from a bag containing 4 white and 7 black balls. If "X" denotes the number of black balls are drawn, then obtain the probability distribution of X. Also find the Mean of the distribution.

222-1st A 323-10000